

## Radon Risk If You Smoke

Radon Level	If 1,000 people who smoked were exposed to this level over a lifetime*...	The risk of cancer from radon exposure compares to**...	WHAT TO DO: Stop smoking and...
<b>20 pCi/L</b>	About 260 people could get lung cancer	250 times the risk of drowning	Fix your home
<b>10 pCi/L</b>	About 150 people could get lung cancer	200 times the risk of dying in a home fire	Fix your home
<b>8 pCi/L</b>	About 120 people could get lung cancer	30 times the risk of dying in a fall	Fix your home
<b>4 pCi/L</b>	About 62 people could get lung cancer	5 times the risk of dying in a car crash	Fix your home
<b>2 pCi/L</b>	About 32 people could get lung cancer	6 times the risk of dying from poison	Consider fixing between 2 and 4 pCi/L
<b>1.3 pCi/L</b>	About 20 people could get lung cancer	(Average indoor radon level)	(Reducing radon levels below 2 pCi/L is difficult.)
<b>0.4 pCi/L</b>	About 3 people could get lung cancer	(Average outdoor radon level)	

Note: If you are a former smoker, your risk may be lower.

**pCi/L** (pico Curies per Liter)

\* Lifetime risk of lung cancer deaths from EPA Assessment of Risks from Radon in Homes (EPA 402-R-03-003).

\*\* Comparison data calculated using the Centers for Disease Control and Prevention's 1999-2001 National Center for Injury Prevention and Control Reports.

This is taken from the EPA website. For more information you may visit:

<http://www.epa.gov/radon/healthrisks.html>